

PROJECTIONS OF CHANGES IN PRODUCTIVITY OF MAJOR AGRICULTURAL CROPS IN THE REPUBLIC OF MOLDOVA ACCORDING TO AN ENSEMBLE FROM 10 GCMS FOR SRES A2, A1B AND B1 EMISSION SCENARIOS IN THE XXI CENTURY

Lilia ȚĂRANU

PhD in Biological Sciences, Senior Consultant in the Area of Adaptation and Vulnerability to Climate Change, Climate Change Office, Ministry of Environment of the Republic of Moldova #156A, Mitropolit Dosoftei St., Office 37, MD 2004, Chişinău, Republic of Moldova E-mails: <u>l.taranu@yahoo.com</u> and <u>clima@mediu.gov.md</u>

Prezentat la 19 august 2013

Rezumat: În articol sunt prezentate rezultatele unui studiu dedicat analizei relației dintre valorile temperaturilor și precipitațiilor medii pe durata perioadei de vegetație și recolta medie la hectar a principalelor culturi agricole la întreprinderile de toate categoriile din Republica Moldova în perioada 1981-2010. S-a stabilit că productivitatea medie la grâul de toamnă, porumbul pentru boabe, floarea soarelui, sfecla de zahăr și tutun, în perioada 1981-2010, a corelat semnificativ din punct de vedere statistic cu valorile temperaturilor și precipitațiilor medii pe durata perioadei de vegetație. Relațiile obținute au fost utilizate pentru estimarea impactului potențial al schimbărilor climatice proiectate de un ansamblu din 10 modele climatice globale impuse de 3 scenarii de emisii ale gazelor cu efect de seră: SRES A2 (înalte), A1B (medii) și B1 (reduse), asupra recoltelor anticipate ale principalelor culturi agricole pentru trei perioade distincte de timp: 2010-2039, 2040-2069 și 2070-2099. În conformitate cu rezultatele obținute, în lipsa măsurilor de adaptare la schimbarea climei, către sfârșitul secolului XXI (perioada temporală 2070-2099), în Republica Moldova se anticipează o reducere semnificativă a productivității principalelor culturi agricole, variind în funcție de scenariul de emisie: între 49 și 74%, la porumbul pentru boabe; 38 și 71%, la grâul de toamnă; 11 și 33%, la floarea soarelui; 10 și 20%, la sfecla de zahăr, respectiv, între 9 și 19%, la tutun. Reducerea semnificativă a productivității principalelor culturi cerealiere se explică prin transferul fazelor fenologice critice în perioade de timp cu condiții nefavorabile de temperatură și umiditate ca urmare a schimbărilor climatice anticipate.

Key words: Crop Yield, Climate Change, Impact Assessment, Statistical Models.

INTRODUCTION

It is now universally accepted that increased atmospheric concentrations of 'greenhouse gases' are the main cause of the ongoing climate change [14] and that these changes are expected to have important effects on different economic sectors (e.g., agriculture, forestry, energy consumptions, tourism, etc.) [16]. Since agricultural practices are climate-dependent and yields vary from year to year depending on climate variability, the agricultural sector is particularly exposed to changes in climate. In Europe, the present climatic trend indicates that in the northern areas, climate change may primarily have positive effects through increases in productivity and in the range of species grown [2], while in southern areas the disadvantages will predominate with lower harvestable yields, higher yield variability and a reduction in suitable areas for traditional crops [37; 9; 35].

For climate change impact assessment, crop growth models have been widely used to evaluate crop responses (development, growth and yield) by combining future climate conditions, obtained from General or Regional Circulation Models (GCMs and RCMs, respectively), with the simulation of CO₂ physiological effects, derived from crop experiments [1]. Many of these impact studies were aimed at assessing crop development shifts and yield variations under changes in mean climate conditions. These analyses showed that increasing temperatures generally shortened the growing period of commercial crops [15], resulting in a shorter time for biomass accumulation. On the other hand, changes in yields were not homogeneous and dependent on crop phenology (e.g. summer and winter crops), crop type (e.g. C and C plants) or environmental conditions (water and nutrient availability) [1; 52; 4; 15; 35]. Other studies stressed that changes in climate variability, as can be expected in a warmer climate, may have a more profound effect on yield than changes in mean climate [40]. Furthermore, the changes in the frequency of extreme climatic events during the more sensitive growth stages have been recognized as a major yield-determining factor for some regions in the future [10; 46]. Temperatures outside the range of those typically expected during the growing season may have severe consequences on crops, and when occurring during key development stages they may have a dramatic impact on final production, even in case of generally favorable weather conditions for the rest of the growing season. Many studies highlighted the potential of heat stresses during the anthesis stage as a yield reducing factor [53; 36; 8], while others pointed out that the joint probability of heat stress-anthesis is likely to increase in future scenarios [33; 47; 46; 2].

Statistical models of crop responses to climate change are the second method for climate change impact assessment in agriculture, based on historical datasets of crop and climate variables have recently been used [20; 21; 22; 44; 9; 41; 42] to address climatic change impacts on agriculture. This method has the advantage of being applicable on large spatial scales, thus facilitating the quantification of impacts in human terms such as levels of risk of hunger [38; 39].

One disadvantage of this method is the potential to introduce significant errors through the linearization of the equations for crop yield [7] and/or the use of monthly data [13], which may not be able to account sufficiently for sub seasonal variability in weather. The validity of empirical methods may be compromised when used with data outside the range for which they were fitted (e.g., climate change).

Mediul

mbiant

This approach can be used for all crops and all regions, but has several deficiencies (as acknowledged by the authors): (i) errors in statistical data of production and in gridded climate data, and (ii) in some instances, low weight of climatic factors for the between year variation in crop yields. Moreover, statistical models cannot be extrapolated without further assumptions to predict production impacts for future conditions (e.g. higher temperature than any historical year, elevated CO₂ concentration [50]. Some authors [28], emphasize three important points that crop models users should consider in future work. First, statistical models represent a very useful, although imperfect tool for projecting climate responses, with all three statistical approaches able to reproduce some of the key aspects of the simulated responses to temperature and precipitation changes. Second, the relative performance of statistical models will depend on the response in question. Time-series models appear particularly good at estimating precipitation responses, while panel or cross-section methods appear more reliable for temperature responses. Finally, the accuracy of statistical approaches depends on the spatial scale of the training data and the scale at which output projections are required. In general, statistical models appear to become more appropriate as the scale of interest becomes broader. It is also at these broader scales that climate projections are most available and reliable, and therefore statistical models are likely to continue to play an important role in anticipating future impacts of climate change [28].

Projecting the qualitative consequences for the Republic of Moldovan agriculture as a result of global climate change is very difficult, because of the uncertainty of variety of natural changes and the lack of reliable modeling tools as crop growth models, which need various input data and a calibration exercise. First assessment of the impact of climate change on crop production in the Republic of Moldova was conducted in the frame of First National Communication of the Republic of Moldova under the UNFCCC [32]. Continuation of researches in this area allowed projecting the effect of direct exposure to elevated CO_2 concentrations in the atmosphere on a yield of winter wheat and maize for different time slices and models [55].

The purpose of our study was: (i) to examine the statistic-empirical relationships between observed mean temperature and precipitation during growing season and average crop yield, based on yield data at the Republic of Moldova's agricultural enterprises of various categories; and (ii) to use these relationships to postulate possible projections of future changes in yield of these crops by 2020's, 2050's and 2080's, based on the projected changes from an ensemble of 10 Global Climate Models (GCM) for three emission scenarios of greenhouse gases and aerosols: SRES A2 (high), A1B (medium) and B1 (low). Typical winter (Triticum aestivum L.) and summer crops (Zea Mays, Helianthus annuus L., Beta vulgaris L. and Nicotiana L.) were considered in this study in order to analyze the specific interactions between the changing climate and crops having different seasonal growth cycles. The interactions phenology-environment in a changing climate were highlighted in order to explain the possible impact on final yield the most vulnerable cereal crops (winter wheat and grain maize).

DATA AND METHODS

The assessment of the climate change impact on agricultural sector was made based on projections of changes in temperature and precipitation received by regionalization of global experiments the most reliable in the Republic of Moldova 10 GCMs for the three SRES A2, A1B and B1 emission scenarios of greenhouse gases and aerosols (see more information in [51]). To assess the vulnerability of main aqricultural crops to climate change was used empirical-statistical approach linking fluctuations of crops vields to climate conditions during the growing season. Statistical analysis of the possible impact of climate change on yield of cereal (winter wheat and grain maize), oil (sunflower) and technical (sugar beet and tobacco) crops was carried out in several steps.

Firstly, according to the statistical data on productivity at agricultural enterprises of various categories were constructed linear and polynomial trends for crop yields in the Republic of Moldova over the two distinct time periods: 1961-1990 (baseline periods) and 1981-2010 (recent periods). Secondly, multiple regression equations linking yield variability with average monthly temperature and precipitation during the agricultural crops growing season, with the highest level of statistical significance were calculated (using the statistical application package STATGRAPH-ICS Centurion and Microsoft Office Excel). The temperature and precipitation predictor variables were selected in conformity with the step by step regression analysis taking into account their contribution to the crops productivity and consecutive analysis of all possible combinations to find the most reliable model. The regression coefficients of the remaining months revealed, in what direction and how much the crops productivity may be modified in response to changes in the temperature and precipitation of the re-

spective month. Finally, the analysis of the impact of future climate changes, determined by temperature and precipitation conditions on the yield of major cereal, oil and technical crops, without undertaken any adaptation measures, was carried out according to a recognized methodological approach [9]. Using the regression equations relationship in the yield variability of major agricultural crops with the temperature and precipitation of the growing season, there were calculated

Mediul

mbiant

projections of future yield changes in the Republic of Moldova, (%/30 years) according to an ensemble from 10 GCMs for the three SRES A2, A1B and B1 emission scenarios, relative to 1981-2010 periods.

RESULTS AND **DISCUS-**SIONS

The vield of most crops generally increased over the past several decades. However, in the most recent decade, yields stagnated for many crops in various regions, whereas temperatures have generally increased. The reasons for this stagnation are debated, and could include agricultural policies [12], climate [23; 5], agronomic and crop management [5] practices, etc.

According to [54], agricultural crops productivity is determined by the level of farming culture, by soil, climatic and weather conditions. Following the generally accepted methodology, we have viewed crop productivity (Yi) as a sum of two elements:

namic average value, determined by the rate of farming intensification and climatic conditions close to average for many years, and the deviation from it ΔY_i (7) is explained by the anomaly of weather conditions of the latter. In other words, the tendencies of crop productivity depend on the implementation of scientific and technical achievements into practice, increased investment into technical means, compliance with the agro technical measures and crop rotation, improved labor organization, better use of fertilizers, modification of varieties used, irrigation, etc. These tendencies are a consequence of gradual improvement in the culture of farming in average soil and climatic conditions.

The linear trends of major aqricultural crops productivity variability on the territory of the Republic of Moldova (RM) for reference period (1961-1990) have been characterized by a sustainable increase of crop yield, by 7.3 g/ha per decade for winter wheat, 3.5 g/ha per decade for grain maize, 0.9 g/ha per decade for sunflower and 25.6 g/ ha per decade for sugar beet (Table 1; Figure 1). Crop productivity increased significantly due to the implementation of intensive technologies and use of irrigation in the 1981-1990 and has reached the maximum level - for winter wheat - 35.0 g/ha; maize for grain - 39.0 g/ha; sunflower - 18.0 g/ha; and sugar beet - 272.5 q/ha.

Then, in the forthcoming three decades (1981-2010) there was a tendency for sharp decrease in crop productivity by 6.2 q/ha per decade for winter wheat, 8.3 g/ha per decade for grain maize, 2.5 g/

Table 1

YIELD LINEAR TRENDS (QUINTAL/YEAR) OF MAJOR CROPS AND THEIR STATISTICAL SIGNIFICANCE (P-VALUE) FOR THE THREE OBSERVATION PERIODS IN THE REPUBLIC OF MOLDOVA

	Yield						
Crop	1961-1990		1981-2010		1991-2010		
	Trend	p-value	Trend	p-value	Trend	p-value	
Winter wheat	0.7288	0.0000	-0.6187	0.0003	-0.6958	0.0297	
Grain maize	0.3480	0.0011	-0.8295	0.0000	-1.0674	0.0042	
Sunflower	0.0862	0.0123	-0.2598	0.0004	-0.0038	0.9697	
Sugar beet	2.5578	0.0090	0.5025	0.6956	0.5025	0.3392	
Tobacco	-	-	-0.0687	0.2486	0.0030	0.9774	

Note: Bold is used to mark statistically significant values.











 $\substack{y=0.0028x^4-0.1427x^3+2.0962x^2-4.6267x+211.58\\R^2=0.3646}$









Figure 1: Yield Variability Trends for the Main Crops and Their Coefficients of Determination (R²) for Two Observation Periods (left - 1961-1990, right - 1981-2010) in the Republic of Moldova

Mediul Amoiant

Table 2

THE INTERRELATIONSHIP OF MAJOR AGRICULTURAL CROPS YIELD VARIABILITY WITH TEMPERA-TURE AND PRECIPITATION DURING THE VEGETATION PERIOD (1981 - 2010)

Сгор	Regression equation	p – value	R ²,%	
Winter wheat	$Y = 130,473 - 2,27358*T_v - 2,60312*T_{v l l} - 0,815868*T_x + 0,0181492*P_{v l} + 0$	0 0003	65.26	
	0,0335445*P _{1X}	0.0005	05.20	
Grain maize	$Y = 177,365 - 2,20195^*T_{v_1} - 1,62369^*T_{v_1} - 3,21222^*T_{v_1} + 0,0169456^*P_{v_2} + 0,016656^*P_{v_2} + 0,01666^*P_{v_2} + 0,0166^*P_{v_2} + 0,016$	0.0002	61 70	
	0,0113687*P _{VII}	0.0002	01.79	
Sunflower	$Y = 41,6803 - 0,817938*T_{VIII} - 0,671806*T_{x} + 0,0242598*P_{VI} - 0,0415422*P_{VIII} - 0,041542*P_{VIII} - 0,04154*P_{VIII} - 0,04154*P_{VIII} - 0,0415*P_{VIII} - 0,0415*P$	0.0011	66.26	
	0,0454977*P _x	0.0011	00.30	
Sugar beet	$Y = 431,527 - 16,9561*T_{v1} + 1,09464*P_{1v} + 0,338911*P_{v} + 0,241995*P_{v1} + 0,241995*P_{v1} + 0,241995*P_{v1} + 0,000$	0.0049	40.02	
	0,314864*P _{v///}	0.0040	49.92	
Tobacco	$Y = 26,6589 - 1,05*T_{y} + 0,0488325*P_{y} - 0,0176172*P_{yyy} - 0,0229145*P_{yy} - 0,0229145*P_{yy} - 0,0229145*P_{yy} - 0,0229145*P_{yy} - 0,0229145*P_{yy} - 0,0176172*P_{yyy} - 0,00229145*P_{yy} - 0,0022914*P_{yy} - 0,0022914*P_{yy}$	0.0004	52.00	
	0,031975*P _x	0.0024	53.20	

Note: Y - yield, quintal/ha; T - average monthly air temperature, ^oC; P - average monthly precipitations, mm; with Roman numerals are noted the corresponding months of vegetation period: since April (IV) to October (X).

ha per decade for sunflower. The greatest decrease in crop productivity was observed in the 1991 - 2010 years, for winter wheat reaching the level of 23.0 q/ha; for grain maize - 19.8 q/ha, while for sunflower - 12.6 q/ha.

The analysis of the data presented in the Table 2 reveals that the influence of climatic conditions during growing season on winter wheat yield in the 1981-2010 periods was statistically significant at 99.9% highest level of significance $(p \leq 0.001)$. Coefficient of determination R² shows that the combined effect of precipitation and temperature defined about 65.3% of the variability of average annual productivity of winter wheat during this period. The regression equation characterizing the interrelation of agricultural crops (corn, sunflower, sugar beet and tobacco) yield variability with the temperature and precipitations during the vegetation period, revealed as well a high level of significance at 99% ($p \le 0.01$). The combined effect of temperature and precipitation during the vegetation period determined the yield variability at the level of 66.4 per cent for sunflower, 61.8 per cent for corn, 53.2 per cent for tobacco and 49.9 per cent for sugar beet.

Crops are known to be sensitive to various aspects of climate. Persistently elevated temperatures have long been known to accelerate progress towards maturity, and more recently have been shown to



Figure 2: Projections of Future Changes in Productivity of Major Agricultural Crops in the Republic of Moldova, (%/30 years) Relative to 1981-2010 Periods, According to an Ensemble from 10 GCMs for SRES A2, A1B and B1 Emission scenarios in the XXI century

have a significant impact on leaf ageing [3; 25]. Crop responses to shorter periods of high temperature, particularly when coincident with flowering, show yield falling dramatically beyond a threshold temperature [31]. Maize is particularly sensitive to hot daytime temperatures, with rapid losses when temperature exceeds 30°C [44; 29]. Crop yields are also sensitive to precipitation. Quantifying the relative effect of temperature and precipitation variability is important for understanding impacts and developing adaptation options for future climatic changes. As temperatures are projected to significantly

increase over the next few decades due to continuing anthropogenic emissions of greenhouse gases, whereas precipitation changes are far less certain [18], this suggests predictability in future crop yields. In addition, it is likely that temperature will have the largest impact as the projected changes are far further outside the range of natural variability than for precipitation changes [24], and because of the seasonal timing of changes in climate [49]. To effectively guide adaptation to future changes, perhaps with different crop growing strategies [43] or selective crop breeding [6], several key questions should

be considered [19]. Firstly, can the relative effects of improved technology, precipitation variability and increasing temperatures be quantified? If so, what is the relative size of the effects of rainfall and hot temperatures on yields? And, what level of technology development may be required to overcome any impact of future climatic changes on yield?

The possible changes in the yield of major agricultural crops (winter wheat, grain maize, sun-flower, sugar beet and tobacco), due to future climate changes in the Republic of Moldova, without undertaken any adaptation measures, is revealed in Figure 2.

As one of the most essential resources to world food supply, wheat yield is very sensitive to temperature change [45; 27]. Lobell & Field (2007) [22], reported about 5.4 % decrease in global mean wheat yield per 1°C increase in temperature, but Asseng et al. (2011) [3] suggested from model simulations that an average growing-season temperature increase by 2°C could cause 50% grain yield reduction in Australia because temperatures more than 34°C stimulate leaf senescence.

Mediul Amolant

The analysis of the obtained results revealed that due to the impact of the main climate indicators (temperature and precipitation) in the RM, productivity of the winter wheat by 2020's could decrease from 9% (SRES A2) to 12% (SRES B1). In comparison with the 1981-2010 time periods, by 2050's the crop productivity may decrease in dependence of the assessed emission scenario from 27% (SRES B1) to 34% (SRES A2). The maximum values of productivity decrease may be reached by 2080's. So, due to changes in values of main climate indicators - precipitation and temperature - the productivity of winter wheat may decrease from 38% (SRES B1) to 71% (SRES A2). The sharp decline in the productivity of winter wheat in the RM can be explained by a shift of critical phenological phases in a more unfavorable period due to temperature increase. The vegetation period of winter wheat (starting with temperatures higher than 5°C in spring), according to an ensemble of 10 GCMs will start in the RM by 2020's earlier by 1 - 4 days (under the SRES A2) and/or by 2 - 6 days (under the SRES B1). By the 2080's, the vegetation period of winter wheat will start earlier by 7-9 days (under the SRES B1) and/ or by 10-13 days (under the SRES A2), in dependence of the assessed emission scenario, with a maximum expected shift in the Central Agro-Ecological Zone (AEZ).

Change of phenology duration is an essential factor for wheat vield. Previous studies had found that warming will shorten wheat phenology duration and decrease wheat yield, mainly due to a shorter growing period, which decreases the duration of photosynthesis and wheat mass accumulation [53; 17]. The reduction would be 4-7% for each 1°C raised [17] and more (see example above). Faroog et al., 2011 [11] have found that wheat yield would decrease with maximum temperatures higher than the optimum temperature in anthesis and grain-filling stages. By use of the index 'sum of effective temperatures above 5°C' there were calculated for winter wheat. according to an ensemble from 10 GCMs for three SRES A2, A1B and B1 emission scenarios, the average initiating date of main development phases in the spring-summer period. Analysis of the data for the most vulnerable Central AEZ pre-

Table 3

PROJECTIONS ON SHIFTING THE PERIOD OF INITIATION OF PHENOLOGICAL PHASES IN WINTER WHEAT DEPENDING ON THE SUM OF EFFECTIVE TEMPERATURES ACCORDING TO AN ENSEMBLE FROM 10 GCMS FOR SRES A2, A1B AND B1 EMISSION SCENARIOS IN THE CENTRAL PART OF THE REPUBLIC OF MOLDOVA

	Winter wheat					
Scenarios	Sum of effective temperatures above 5°C necessary to initiate the phenological phase					
	Tiller initiating	Jointing	Kernel in milk	Kernel in dough		
	(125ºC)	(455⁰C)	(685ºC)	(945ºC)		
1961-1990	24/04	27/05	13/06	30/06		
2020's						
A2	20/04	22/05	07/06	24/06		
A1B	18/04	20/05	06/06	23/06		
B1	18/04	20/05	06/06	23/06		
2050's						
A2	15/04	17/05	02/06	18/06		
A1B	13/04	16/05	31/05	17/6		
B1	15/04	17/05	03/06	19/06		
2080's						
A2	10/04	12/05	28/05	12/06		
A1B	09/04	10/05	26/04	11/04		
B1	14/04	15/05	31/04	18/06		

Mediul Annoiant

Table 4

PROJECTIONS ON SHIFTING THE PERIOD OF INITIATION OF PHENOLOGICAL PHASES IN MAIZE VARIETIES OF DIFFERENT MATURITY GROUPS DEPENDING ON THE SUM OF EFFECTIVE TEMPERA-TURES ACCORDING TO AN ENSEMBLE FROM 10 GCMS FOR SRES A2, A1B AND B1 EMISSION SCENAR-IOS IN THE CENTRAL PART OF THE REPUBLIC OF MOLDOVA

	Maize varieties of different maturity groups (early, intermediate and late)					
Sacaria	Sum of effective temperatures above 10°C necessary to initiate the phenological phase					
Scenario	Germination - tasseling					
	(320-410°C)	(400-510°C)	(520-660°C)			
1961-1990	23/06	02/07	16/07			
2020's						
A2	16/06	25/06	08/07			
A1B	14/06	23/06	05/07			
B1	14/06	23/06	05/07			
		2050's				
A2	09/06	18/06	29/06			
A1B	06/06	15/06	27/06			
B1	09/06	18/06	03/07			
	2	2080's				
A2	01/06	09/06	20/06			
A1B	31/05	08/06	20/06			
B1	06/06	15/06	27/06			
Scenario		Sowing – milky grain				
10(1,1000	(720-770°C)	(820-870°C)	(880-930°C)			
1961-1990	26/07	04/08	09/08			
4.2	17/07	2020's	20/07			
AZ	1//0/	24/07	29/07			
AIB	14/07	21/07	26/07			
BI	14/07	22/07	27/07			
4.2	08/07	2050's	10/07			
A2	08/07	15//	19/07			
AIB	05/07	12/07	16/07			
DI	09/07	10/07	21/07			
<u> </u>	28/06	04/07	08/07			
A1R	28/06	05/07	11/07			
R1	05/07	12/07	17/07			
BI	05/07	Sowing – milky dough grain	17/07			
Scenario	(770-820°C)	(870-920°C)	(970-1020°C)			
1961-1990	31/07	08/08	17/08			
		2020's	11100			
A2	20/07	28/07	05/08			
A1B	18/07	26/07	02/08			
B1	18/07	26/07	03/08			
		2050's				
A2	11/07	18/07	25/07			
A1B	09/07	15/07	22/07			
B1	13/07	20/07	27/07			
2080's						
A2	01/07	08/07	14/07			
A1B	01/07	08/07	15/07			
B1	09/07	16/07	23/07			
	07/07	10/07	25/07			

sented in the table below, revealed that by 2020's output in the tiller initiating phase at winter wheat may have shifted in average from 4 days (SRES A2) up to 6 days (SRES B1) (Table 3). By 2050's the shift in the respective phenological phases will account from 9 days (SRES A2) up to 11 days (SRES A1B), while by 2080's according to the SRES

A2 μ A1B emission scenarios the tiller initiating phase will start by 14 days early the onset of respective phase in the reference period (1961-1990).

According to the projections, by 2020's the phenological phase of jointing may come for winter wheat earlier from 4 days (SRES A2), up to 7 days (SRES B1). According to the performed assessment, the humidity conditions in this period will be close to optimal (HTC=1.0), according to L. Taranu (2013) (unpublished data). However, by 2080's this shift can already draw from 12 days (SRES B1), up to 15-17 days (SRES A2 and A1B). Humidity conditions for this period would be sufficient only in accordance with the SRES B1 low emission scenario (HTC=1.0), while according the other two emission scenarios (SRES A2 and A1B) there will be recorded insufficient humidity conditions (HTC=0.8-0.9), thus the critical period for jointing at winter wheat will take place in dryer conditions, which will impact a sharp decrease in the productivity.

Lobell & Asner 2003 [23], evaluated maize and soybean production relative to climatic variation in the United States, reporting a 17% reduction in yield for every 1°C rise in temperature, but this response is unlikely because the confounding effect of rainfall was not considered. In a recent evaluation of global maize production response to both temperature and rainfall over the period 1961-2002 [22] reported an 8.3% yield reduction per 1°C rise in temperature. The analysis of obtained results revealed, that due to changes in heat and water regime during the growing season, yield of grain maize in RM by 2020's may decrease from 9-14% (SRES A2 and B1) to 16% (SRES A1B). In comparison with the 1981-2010 years, by 2050's the crop productivity may decrease in dependence of the assessed emission scenario from 35% (SRES B1) to 48-52% (SRES A2 and A1B). The maximum values of productivity decrease may be reached by 2080's. So, due to changes in values of main climate predictor variables – precipitation and temperature - the productivity of grain maize may decrease from 49% (SRES B1) to 74% (SRES A1B). By use of the index 'sum of

<u>Mediul</u> Amplant

effective temperatures above 10°C' there were calculated for maize of different maturity groups, according to an ensemble from 10 GCMs for three SRES A2, A1B and B1 emission scenarios, the possible average initiating date of main development phases in the springsummer period. Analysis of the data for the most vulnerable Central AEZ presented in Table 4, revealed that by 2020's onset in the germination tasseling phase at maize varieties may have shifted, in dependence from emission scenario and maturity group from 7 days (SRES A2) up to 9 days (SRES B1) for the early and intermediate; and from 8 days (SRES A2) up to 11 days (SRES B1) for late maturity groups. While by 2080's, according to SRES B1 и A2 emission scenarios, the germination - tasseling phase will start by 17-22 days (for the early and intermediate); and by 20-26 days (for late maturity groups) early the onset of respective phase in the reference period (1961-1990) (Table 4). In this regard, it is interesting to note that the actual observed changes in the occurrence of phenological germination - tasseling phase of maize due to the temperature increase over the last two decades (1991-2010) were by 5 days, for early and intermediate, and by 6 days, for late maturing hybrids.

According to the projections, by 2020's onset in the sowing – milky grain phase at maize hybrids may have shifted, in dependence from emission scenario and maturity group from 9 days (SRES A2) up to 12 days (SRES B1) for the early maturity group; and from 11 days (SRES A2) up to 14 days (SRES B1) for intermediate and late maturity groups. While by 2080's, according to the SRES B1 и A2 emission scenarios, the sowing - milky grain phase will start early the onset of respective phase in the 1961-1990 period by 21-30 days for the early maturity group; and by 23-32 days for late maturity group (Table 4). The actual observed changes in the occurrence of phenological sowing milky grain phase at maize due to the temperature increase over past two decades (1991-2010) were by 7 days for early maturity group and by 9 days for intermediate and late maturing hybrids. According to the projections, in the 2020's the humidity conditions will be close to optimal (HTC=1.2) during the sowing – tasseling phenological phase across all three scenarios for early and intermediate hybrids, while for late maturity corn hybrids moisture conditions will be sufficient only under two emission scenarios: SRES A2 and B1 (HTC = 1.0).

The next phenological stage the flowering - milky ripening at maize will be ongoing in low humidity conditions (HTC=0.9-1.0). By the 2080's the humidity conditions would be close to optimal (HTC=1.0) during the tasseling phenological stage under SRES A1B and B1 emission scenarios; and insufficient (HTC=0.8) for the SRES A2 high emission scenario. The subsequent critical phenological phases as flowering and milk grain ripening at maize regardless of maturity group will take place in dryer conditions according to high emission SRES A2 (HTC=0.6) and medium emission SRES A1B (HTC=0.8) scenarios; or in insufficient humidity conditions (HTC=0.9) under the low emission scenario SRES B1, which along with an increase of temperatures will reduce dramatically the crop yield.

By the end of the XXI century, the cultivation of major cereal crops such as the winter wheat and maize will be either impossible according to the SRES A2 high emission scenario or economically not cost effective according to the SRES A1B medium and SRES B1 low emission scenarios (Figure 2), due to changes in climatic conditions, without adaptation measures (*if maintaining the current cultivation technologies and used varieties*) in the Republic of Moldova.

For oil crop such as sunflower, which is relatively drought-resistant, there are projected more favorable climate conditions during the growing season than for cereal crops: winter wheat and grain maize. Slight increase in productivity by 2-4% it is possible for sunflower in the 2020's, which will be replaced in the future with consecutive decline in yield. By 2050's, the most severe decrease in productivity by 10-13% is projected under the SRES A2 high and SRES A1B medium emission scenario. While under the SRES B1 low emission scenario the projection is more favorable, it is expected a decrease of productivity only by 7%. In 2080's, the most severe yield reduction for sunflower in the RM will be observed according to SRES A2 high emission scenario - by 33%, while for SRES B1 low emission scenario the forecast is more favorable - a decrease by 11% (Figure 2).

For sugar beet by 2020's, when assessing the combined effect of temperature and precipitation during the growing season, it is expected a decrease in productivity by 1-5% under two assessed emission scenarios SRES B1 and A1B. By 2050's, there will persisting the decreasing trend in productivity due to climate changes. The most severe decrease in productivity, by 10-12% is predicted under the SRES A1B medium and A2 high emission scenario. While under the SRES B1 low emission scenario the projection is more favorable, it is predicted a decrease of productivity only by 5%. In 2080's, the most severe yield reduction for sugar beet in the RM will be observed according to SRES A2 high emission scenario - by 20%, while for SRES B1 low emission scenario the forecast is more favorable – a decrease by 10% (Figure 2).

For other high added value technical crop – tobacco, similar changes are expected: a slight decrease in the yield by 1%, according to the SRES A2 and A1B scenarios in the 2020's. By 2050's, there will persisting the decreasing trend in productivity due to climate changes. The most severe decrease in productivity, by 8-10% is predicted under the SRES A1B medium and SRES A2 high emission scenarios. While under the SRES B1 low emission scenario the projection is more favorable, it is predicted a decrease of productivity only by 6%. In 2080's, the most severe yield reduction for sugar beet in the RM will be observed according to SRES A2 high emission scenario – by 19%, while for SRES B1 low emission scenario the forecast is more favorable – a decrease by 9% (Figure 2).

In this paper, relationships between observed mean temperature and precipitation during growing season and average crop yield based on statistical data at the Republic of Moldova's agricultural enterprises of various categories are explored and then used to estimate potential impacts of climate change scenarios on anticipated average yields for the 2020's, 2050's and 2080's time periods. Average yields of winter wheat, grain maize, sunflower, sugar beet and tobacco in the 1981-2010 years were highly correlated with precipitation and temperature during the growing period. Based on range of available temperature and precipitation climatic conditions during the vegetation period projected by an ensemble from 10 GCMs for SRES A2 (high), A1B (medium) and B1 (low) emissions scenarios average agricultural crop yields achievable in the Republic of Moldova could decrease by about 59 to 87% for grain maize; 38 to 71% for winter wheat; 11 to 33% for sunflower, 10 to 20% for sugar beet; and 9 to 19% for tobacco by 2080's, not including the direct effect of increased atmospheric CO₂ concentration, advances in plant breading and crop production practices or changes in the impacts of weeds, insects and diseases on yield. The sharp decline in the productivity of cereal crops can be explained by a shift of critical phenological phases in the more unfavorable humidity and temperature conditions due to climate change.

CONCLUSIONS

The impact assessment performed allow conclude that the negative effect of global warming, according to an ensemble from 10 GCMs for SRES A2 (high), A1B (medium) and B1(low) emission scenarios in the XXI century will not be offset by increase of precipitations. In these circumstances, without undertaken any adaptation measures it can be expected by 2080's: a significant drop in the productivity for grain maize from 49% (SRES B1) to 74% (SRES A1B) and winter wheat from 38% (SRES B1) to 71% (SRES A2); a medium drop in the productivity for sunflower from 11% (SRES B1) to 33% (SRES A2), respectively for sugar beet from 10% (SRES B1) to 20% (SRES A2); and for tobacco from 9% (SRES B1) to 19% (SRES A2), in comparison with the average productivity of the Republic of Moldova's major agricultural crops in the most recent period of 1981-2010.

BIBLIOGRAPHY

1. Ainsworth E.A., Long S.P. (2005): What have we learned from 15 years of free-air CO_2 enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO_2 . New Phytol, 165:351–372

2. Alcamo J., Moreno J.M., Nováky B. et al (2007): Europe. Climate change 2007: impacts, adaptation and vulnerability. In: Parry M.L. et al (eds) Contribution of working group II to the fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge, pp 541–580

3. Asseng S., Foster I. T., and Turner N. C. (2011): The impact of temperature variability on wheat yields. Glob. Change Biol. 17:997– 1012.

4. Brassard J.P., Sigh P. (2008): Impact of climate change and CO_2 increase on agricultural production and adaptation options for Southern Quebec, Canada. Mitig Adapt Strategies Glob Change, Vol. 13:241-265.

5. Brisson N., Gate P., Gouache D., Charmet G., Oury F.-X., Huard F. (2010): Why are wheat yields stagnating in Europe? A comprehensive data analysis for France. *Field Crops Research*, 119,

<u>LuibeW</u> Anclount

201–212. Calderini D, Slafer G. (1998): Changes in yield and yield stability in wheat during the 20th century. *Field Crops Research*, 57, 335–347.

6. Cattivelli L., Rizza F., Badeck F.-W. et al. (2008): Drought tolerance improvement in crop plants: an integrated view from breeding to genomics. *Field Crops Research*, 105, 1–14. doi: 10.1016/j.fcr.2007.07.004.

7. Challinor A. J., Wheeler T. R, Osborne T. M., and Slingo J. M. (2006): Assessing the vulnerability of crop productivity to climate change thresholds using an integrated crop–climate model. *Avoiding Dangerous Climate Change*, J. Schellnhuber et al., Eds, Cambridge University Press, 187–194.

8. Challinor, A. J., Wheeler T. R., Slingo J. M. (2005): Simulation of the impact of high temperature stress on the yield of an annual crop. *Agric For Meteorol*, Vol. 135: 180–189.

9. Constantinov T., Taranu L., Mironova T., Toderas V. (2009): Assessment of vulnerability of some agricultural crops productivity in the Republic of Moldova, without adaptation measures, to new climate conditions according to the CSIRO-Mk2, HADCM2 and ECHAM4 general atmospheric models. The Bulletin of the Academy of Sciences of Moldova, Series of "Biological, Chemical and Agricultural Sciences". Republic of Moldova, Chisinau, 2009, Nr.2 (308). C. 124-132.

10. Easterling W., Apps M. (2005): Assessing the consequences of climate change for food and forest resources: a view from the IPCC. *Climate Change*, Vol. 70: 165–189.

11. Farooq M., H. Bramley J. A. Palta, and K. H. M. Siddique. (2011): Heat Stress in Wheat during Reproductive and Grain-Filling Phases. Crit. Rev. Plant Sci. 30:491–507.

12. Finger R (2010): Evidence of slowing yield growth - the example of Swiss cereal yields. Food Policy, 35, 175–182. doi: 10.1016/j. foodpol.2009.11.004.

13. Fischer, G., M. Shah, F. N. Tubiello, and H. van Velhuizen (2005): Socio-economic and climate change impacts on agriculture: An integrated assessment, 1990–2080. *Philos. Trans. Roy. Soc.*, 1463, 2067–2083.

14. Forster, P., V. Ramaswamy, P. Artaxo et al. (2007): Changes in atmospheric constituents and in radiative forcing. *In*: Solomon S., et al (eds.) Climate change 2007: the physical science basis. Contribution of working group I to the fourth assessment report of the intergovernmental panel on climate change. Cambridge University Press, Cambridge.

15. Giannakopoulos C., P. Le Sager, M. Bindi et al. (2009): Climatic changes and associated impacts in the Mediterranean resulting from global warming. *Global Planet Change*, Vol. 68: 209–224.

16. Hanson, C. E., J. P. Palutikof, M. T. J. Livermore et al. (2007): Modeling the impact of climate extremes: an overview of the MICE Project. *Climate Change*, Vol. 81:163–177.

17. Hatfield, J. L., K. J. Boote, B. A. Kimball et al. (2011): Climate impacts on agriculture: implications for crop production. Agron. J. 103:351–370.

18. Hawkins E., Sutton R. (2011): The potential to narrow uncertainty in projections of regional precipitation change. *Climate Dynamics*, 37, 407–418. doi: 10.1007/s00382-010-0810-6.

19. Hawkins E., Fricker T. E., Challinor A. J. et al. (2013): Increasing influence of heat stress on French maize yields from the 1960s to the 2030s. *Global Change Biology* 19, 937–947, doi: 10.1111/ gcb.12069

20. Iglesias A., Quiroga S. (2007): Measuring the risk of climate variability to cereal production at five sites in Spain. *Clim.Res.* 34:47–57

21. Iglesias A., C. Rosenzweig and D. Pereira (2000): Agricultural impacts of climate change in Spain: Developing tools for a spatial analysis. *Global Environ. Change*, 10, 69–80. 22. Lobell D. and Field C. (2007): Global scale climate–crop yield relationships and the impacts of recent warming. *Environ. Res. Lett.* 2 014002 doi:10.1088/1748-9326/2/1/014002

23. Lobell D.B., Asner G.P. (2003): Climate and management contributions to recent trends in U.S agricultural yields. *Science*, 299, 1032, doi: 10.1126/science.1078475.

24. Lobell D.B., Burke M.B. (2008) Why are agricultural impacts of climate change so uncertain? The importance of temperature relative to precipitation. *Environmental Research Letters*, 3 (034), 007. doi: 10.1088/1748-9326/3/3/034007.

25. Lobell D.B., Sibley A., Ortiz-Monasterio J.I. (2012): Extreme heat effects on wheat senescence in India. *Nature Climate Change*, 2, 186–189. doi: 10.1038/ nclimate1356.

26. Lobell D. B., M. B. Burke, C. Tebaldi et al. (2008): Prioritizing climate change adaptation needs for food security in 2030. *Science*, 319, 607–610.

27. Lobell D. B., W. Schlenker, and J. Costa-Roberts. (2011): Climate Trends and Global Crop Production since 1980. *Science* 333:616–620.

28. Lobell, D.B., Burke, M.B. (2010): On the use of statistical models to predict crop yield responses to climate change. *Agric. Forest Meteorol.* doi:10.1016/j. agrformet.2010.07.008

29. Lobell D.B., M. Banziger, C. Magorokosho, and B. Vivek. (2011a): Nonlinear heat effects on African maize as evidenced by historical yield trials. *Nature Climate Change* 1(1): 42-45.

30. Lobell D.B., W.S. Schlenker, and J. Costa-Roberts. (2011b): Climate trends and global crop production since 1980. *Science* 333(6042): 616-620.

31. Luo Q. (2011): Temperature thresholds and crop production: a review. *Climatic Change*, 109, 583– 598. Meehl GA, Stocker TF, Collins W et al. (2007): Global climate projections. In: Climate Change 2007: The Physical Science Basis.



Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change (eds: Solomon S. et al.). Cambridge University Press, UK.

32. Ministerul Mediului și Amenajării Teritoriului / PNUD Moldova (2000), Prima Comunicare Națională a Republicii Moldova elaborată în cadrul Convenției Națiunilor Unite privind Schimbarea Climei., Chișinău. – 2000. - 74 p.

33. Moriondo M., Bindi M. (2006): Comparison of temperatures simulated byGCMs, RCMs and statistical downscaling: potential application in studies of future crop development. *Clim Res* 30:149–160

34. Moriondo M., Giannakopoulos C., Bindi M. (2011): Climate change impact assessment: the role of climate extremes in crop yield simulation. *Climatic Change*. 104:679–701DOI 10.1007/s10584-010-9871-0

35. Moriondo M., Giannakopoulos C., Bindi M. (2010): Climate change impact assessment: the role of climate extremes in crop yield simulation. *Climatic Change*, doi 10.1007/s10584-010-9871-0.

36. Morrison M.J, Stewart D.W. (2002): Heat stress during flowering in summer brassica. Crop Sci 42: 797–803

37. Olesen J. E. and 16 co-authors (2007): Uncertainties in projected impacts of climate change on European agriculture and terrestrial ecosystems based on scenarios from regional climate models. *Climatic Change*: 81:123–143.

38. Parry M., C. Rosenzweig, A. Iglesias et al. (2004): Effects of climate change on global food production under RSES emissions and socio-economic scenarios. *Global Environ. Change*, 14, 53–67.

39. Parry M., C. Rosenzweig and M. Livermore (2005): Climate change, global food supply and risk of hunger. *Philos. Trans. Roy. Soc.*, 360, 2125–2138.

40. Porter J.R., Semenov M.A. (2005): Crop responses to climatic variation. Philos Trans R Soc Lond B Biol Sci 360:2021–2035 41. Quiroga S., Iglesias A. (2009): A comparison of the climate risks of cereal, citrus, grapevine and olive production in Spain. *Agric Syst.* 101:91–100.

42. Quiroga S, Iglesias A. (2010): A comparison of the climate marginal productivity on representative crop yields in Spain. *Princip: Estud Econ Política*. 16:55–70.

43. Rosenzweig C., Tubiello F. (2007): Adaptation and mitigation strategies in agriculture: an analysis of potential synergies. Mitigation and Adaptation Strategies for Global Change, 12, 855–873. doi: 10.1007/s11027-007-9103-8.

44. Schlenker W., Roberts M.J. (2009): Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change. *PNAS USA*, 106, 15594–15598, doi:10.1073/ pnas.0906865106.

45. Schmidhuber J. and F. N. Tubiello (2007): Global food security under climate change. *Proc. Natl Acad. Sci. USA* 104:19703–19708.

46. Schneider. S. H., S. Semenov; A. Patwardhan et al. (2007): Assessing key vulnerabilities and the risk from climate change. In: Parry ML et al. (eds.) Climate change 2007: impacts, adaptation and vulnerability. Contribution of working group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, pp 779-810.

47. Semenov M.A. (2007): Development of high-resolution UKCIP02-based climate change scenarios in the UK. *Agric For Meteor.* 144:127–138.

48. Semenov M.A., Barrow E.M. (1997): Use of a stochastic weather generator in the development of climate change scenarios. *Clim Change*. 35:397–414.

49. Semenov M.A., Shewry P.R. (2011): Modelling predicts that heat stress, not drought, will increase vulnerability of wheat in Europe. *Science Report*, 1, 66. doi: 10.1038/srep00066.

50. Soussana J., Graux A. and Tubiello F. N. (2010): Improving

the use of modeling for projections of climate change impacts on crops and pastures. *Journal of Experimental Botany*, Vol. 61, No. 8, pp. 2217–2228.

51. Taranu L., Bercu I., Deveatii D. (2012): Regional Climate Change Scenarios for the Republic of Moldova: Future Temperature and Precipitation Projections from Ensembles of 10 Global Climate Models. *Mediul Ambiant*. Nr.3 (63), P. 33-42.

52. Thomson A.M., Brown R.A., Rosenberg N.J. et al. (2005): Climate change impacts for the conterminous USA: an integrated assessment—part 3 dry land production of grain and forage crops. *Clim Change*. 69:43–65.

53. Wheeler T.R., Craufurd P.Q., Ellis R.H. et al. (2000): Temperature variability and the yield of annual crops. *Agric Eco Environ*. 82:159–167.

54. Дарадур М. И. (2001): Изменчивость и оценки риска экстремальных условий увлажнения. 160 р.

55. Коробов Р., Чалык С., Буюкли П. (2004): Оценка чувствительности растениеводства к возможному изменению климата. В кн.: Климат Молдовы в XXI веке: проекции изменений, воздействий, откликов. Кишинев. стр. 213 – 253.